



Atlas of Pancreaticobiliary Maljunction & Congenital Biliary Dilatation



Japanese Study Group on Pancreaticobiliary Maljunction (JSGPBM)
Scientific Committee

Editor-in-Chief, Kenitiro Kaneko
Shin Ishihara
Shigeru Ono
Hironao Miyoshi
Yoshiki Morotomi

In collaboration with Naoto Urushihara

Preface

Pancreaticobiliary maljunction (PBM) was discovered as a malformation accompanying choledochal cysts. In Japan, where choledochal cysts are common, the concept of PBM is widespread, but it seems unfamiliar in other countries where choledochal cysts are rare. However, the term “choledochal cyst” includes various entities of congenitally dilated bile ducts with or without PBM. Almost all choledochal cysts seen in Asia have localized dilatation of the bile duct and PBM, where PBM plays a major role in pathophysiology. In countries outside Asia, the frequency of choledochal cysts without PBM is relatively high, leading to confusion when discussing “choledochal cysts”. We felt it necessary to differentiate this type with PBM from other types without PBM and defined the

common type with PBM in Asia as “congenital biliary dilatation”.

PBM is also observed in patients without bile duct dilatation. Gallbladder cancers arise at an extraordinarily high frequency in these non-dilated PBM patients. However, the definition of non-dilatation has been unclear. We have developed ultrasonographic criteria for bile ducts of normal diameter, which we present here.

In this Atlas, we attempt to demonstrate clear and precise explanations of PBM and congenital biliary dilatation and hope this Atlas will help people unfamiliar to this field. The contents of the Atlas are free to use providing the source (JSGPBM Atlas of PBM & CBD) is cited.

September 2022
Editor-In-Chief, Kenitiro Kaneko

Contents

Introduction

- What is PBM? 6–8
- What is CBD? 9–10
- Localized BD dilatation 11
- Embryogenesis 12
- Treatment of PBM 13
- Biliary carcinogenesis 14–15
- Surgery for CBD 16
- Recommended terms 17

PBM, pancreaticobiliary maljunction
CBD, congenital biliary dilatation
BD, bile duct

Basics

- Pathophysiology, diagnosis & classification 18–38
- What is CBD? 39–46
- Typical CBD 47–56
- Nondilated PBM 57–67
- Treatment of PBM 68–83
- Postoperative complications 84–94
- High confluence of the pancreaticobiliary ducts 95–97

Advanced

- Issues on Todani's classification 98–112
- Nondilated type of PBM
 - Definition, pathophysiology, and diagnosis 113–123
- PBM seemingly complicated & rare type 124–128
- Common channel & narrow segment 129–138
- What causes biliary dilatation? 139–146
- Pathophysiology in pediatric patients
 - Mechanism of protein plug formation 147–157
 - Symptomatology besides protein plug 158–162
- Pathophysiology in adult patients
- Bile duct cancer in nondilated PBM
- Not typical cases
- Pancreas divisum and PBM
- Treatment of type IV-A

Expert

- Prenatal diagnosis,
Transitional type to Biliary Atresia 163–174
- Biliary carcinogenesis during childhood
- Double bile ducts and PBM
- Rare types of choledochal cysts
 - Type Ib 160
 - Type II, diverticulum
 - Type III, choledochoceles
 - Type IV-B really exists?
 - Type V, Caroli syndrome & Caroli disease

Introduction


First version, September 2022

What is **Pancreaticobiliary Maljunction (PBM)**?

Definition

A congenital malformation in which the **pancreatic duct** and **bile duct** join **outside the duodenal wall**

The junction outside of the sphincter action is a functional phenomenon of PBM but is not the definition of it. However, when the confluence outside the duodenal wall cannot be shown, identification of the junction outside the sphincter action is enough for a diagnosis of PBM.
See Basic Section.




What is **Pancreaticobiliary Maljunction (PBM)**?

Pathophysiology

In PBM, the sphincter action does not regulate the pancreaticobiliary junction and **two-way regurgitation** of pancreatic juice and bile occurs, resulting in various pathological conditions of the biliary tract and pancreas.

During **childhood**, intermittent **symptoms** including abdominal pain, vomiting, and jaundice occur. During **adulthood**, **biliary cancers** develop.



Normal


PBM

What is **Pancreaticobiliary Maljunction?**

Two major categories according to whether the bile duct is dilated or not:

- Dilated type = congenital biliary dilatation (CBD)
- Non-dilated type

The definition of non-dilated type remains problematic.
See *Basic and Advanced Sections*.
Existence of non-dilated type denies that
pancreatobiliary regurgitation causes biliary dilatation.
See *Advanced Sections*.




What is Congenital Biliary Dilatation (CBD)?

Definition


Congenital malformation involving **PBM** and **localized dilatation of the bile duct (BD)** including the common bile duct

Definition of CBD proposed by the Japanese Study Group on Pancreaticobiliary Maljunction (JSGPBM) in 2015



In this case, the common hepatic duct (CHD) and the base of the cystic duct (CD) are dilated, but bile ducts upstream to the CHD and the gallbladder are not dilated (localized dilatation).


The term choledochal cyst involves different entities



“Localized dilatation of BD” indicates “congenital” disease


Congenital biliary dilatation

Even when dilatation extends to intrahepatic BDs, dilated areas are localized. **Dilated BDs abruptly change to those of normal caliber (→).** The GB is never enlarged, although the cystic duct is dilated at its base. However, in secondary dilatation, dilatation extends smoothly upstream to the peripheral intrahepatic BD. GB is also enlarged (Courvoisier sign).



Protein plug
compacted in the common channel


Secondary dilatation due to obstruction by BD carcinoma



Smoothly dilated until peripheral intrahepatic BD.
GB is also enlarged.

PBM arises from anomalous genesis of ventral pancreatic duct and bile duct


What is wrong in this figure?



Answer: Santorini's duct (dorsal PD) is depicted behind the CBD. It must be ventral to the CBD.

Pancreas development

Ventral pancreatic bud rotates behind the duodenum and fuses with dorsal bud



Connection anomaly never happens between the BD and dorsal pancreatic duct in PBM. Fusion failure between the ventral and dorsal pancreatic ducts (i.e., pancreas divisum) can occur together with PBM. See *Advanced Section*.

Treatment for pancreaticobiliary maljunction

- Reasons for treatment necessity

PBM causes:

Not merely to mend malformation!


- ✓ Recurrent **symptoms** including abdominal pain, vomiting, and jaundice in children
- ✓ High occurrence of **biliary cancers** in adults see next

- **Aim** of treatment = To **prevent symptoms and cancers**
- The following must be undertaken:
 - **Bile duct resection** for congenital biliary dilatation
 - **At least cholecystectomy** for non-dilated PBM*

*No consensus on the need for bile duct resection because little incidence of carcinogenesis at the BD

Biliary carcinogenesis


1,511 adult cases of PBM
Japan nationwide survey 1990–2007



Cancer rate

CBD 21.6%

215/997 cases



Non-dilated PBM 42.4%

218/514 cases


Biliary cancers arise 10–20 years earlier in PBM

Gallbladder cancer			P
PBM	Yes	No	
	36 cases	232 cases	
Age (years)	58.9 ± 8.8	69.4 ± 8.7	<0.01
M:F	1:2.1	1:2.9	NS

Bile duct cancer			P
PBM	Yes	No	
	6 cases	159 cases	
Age (years)	54.8 ± 14.3	68.8 ± 9.1	<0.01
M:F	1:2	1:0.7	NS


Kamisawa T, Tu Y, Kuwata G, Egawa N, Nakajima H, Tsuruta K, Okamoto A, Matsukawa M. Biliary carcinoma risk in patients with pancreaticobiliary maljunction and the degree of extrahepatic bile duct dilatation. Hepatogastroenterology. 2006 Nov-Dec;53(72):816-8.

Surgery for CBD



NS, narrow segment

Must resect where cancers would arise and stop the reflux of pancreatic juice to prevent cancers and symptoms



Recommended terms

- **Pancreaticobiliary maljunction (PBM)**
- **Congenital biliary dilatation (CBD)**
- **Bile duct resection** as a procedure for CBD

Non-recommended terms

- Pancreaticobiliary malunion or pancreatobiliary malunion
- Choledochal cyst (when indicating CBD (Todani Ia, c, and IV-A))

Choledochal cysts involve different entities such as CBD, choledochoceles, and Caroli's disease.